



Your Discoveries  
Begin with US.

Search:

[Home](#) | [Ordering Info](#) | [Technical Help](#) | [About ATCC](#) | [Contact Us](#)

## Search

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's [Material Transfer Agreement](#).

Customers in Europe, Australia, Japan, Korea, New Zealand and Taiwan must contact a [local distributor](#) for pricing information and to place an order for ATCC cultures and products.

Cell Lines	
<b>ATCC Number:</b> CRL-1651	<b>Price:</b> \$175.00
<input type="button" value="Order this item"/>	
<b>Designation:</b> COS-7	<b>Depositors:</b> Y. Gluzman
<b>Biosafety Level:</b> 2	<b>Shipped:</b> frozen
<b>Medium &amp; Serum:</b> <a href="#">See Propagation</a>	<b>Growth Properties:</b> adherent
<b>Organism:</b> <i>Cercopithecus aethiops</i> (African green monkey)	<b>Morphology:</b> fibroblast
<b>Tissue:</b> kidney; SV40 transformed	
<b>Cellular Products:</b> T antigen	
<b>Permits/Forms:</b> In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please <a href="#">click here</a> for information regarding the specific requirements for shipment to your location.	
<b><a href="#">Related Cell Culture Products</a></b>	
<b>Comments:</b>	This line was derived from the CV-1 cell line (ATCC CCL-70) by transformation with an origin defective mutant of SV40 which codes for wild-type T antigen. [1822] The sale of any endangered wildlife is conditional on the issuance of a valid permit by the U.S. Fish and Wildlife Service and will not be consummated until a permit is obtained from the U.S. Fish and Wildlife Service.
<b>Applications:</b>	transfection host
<b>Virus Susceptibility:</b>	SV40 (lytic growth); SV40 tsA209 at 40C; SV40 mutants with deletions in the early region
<b>Propagation:</b>	ATCC medium: Dulbecco's modified Eagle's medium with 4 mM L-glutamine adjusted to contain 1.5 g/L sodium bicarbonate and 4.5 g/L glucose, 90%; fetal bovine serum, 10% Temperature: 37.0 C

<b>Subculturing:</b>	Remove medium, and rinse with 0.25% trypsin, 0.03% EDTA solution. Remove the solution and add an additional 1 to 2 ml of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37C) until the cells detach. Add fresh culture medium, aspirate and dispense into new culture flasks.
<b>Split Ratio:</b>	A subcultivation ratio of 1:4 to 1:8 is recommended
<b>Fluid Renewal:</b>	2 to 3 times per week
<b>Freeze Medium:</b>	culture medium 95%; DMSO, 5%
<b>Related Products:</b>	Recommended medium (without the additional supplements or serum described under ATCC Medium) - ATCC 30-2002 recommended serum - ATCC 30-2020 parental cell line - ATCC CCL-70
<b>References:</b>	<p><u>1822</u>: Gluzman Y. SV40-transformed simian cells support the replication of early SV40 mutants. Cell 23: 175-182, 1981. PubMed: <u>6260373</u></p> <p><u>32447</u>: Fernandez LM, Puett D . Lys583 in the third extracellular loop of the lutropin/choriogonadotropin receptor is critical for signaling. J. Biol. Chem. 271: 925-930, 1996. PubMed: <u>8557706</u></p> <p><u>32459</u>: Maestrini E, et al . A family of transmembrane proteins with homology to the MET-hepatocyte growth factor receptor. Proc. Natl. Acad. Sci. USA 93: 674-678, 1996. PubMed: <u>8570614</u></p> <p><u>32500</u>: Campbell M, et al . The simian foamy virus type 1 transcriptional transactivator (Tas) binds and activates an enhancer element in the gag gene. J. Virol. 70: 6847-6855, 1996. PubMed: <u>8794326</u></p> <p><u>32502</u>: Gonzalez Armas JC, et al . DNA immunization confers protection against murine cytomegalovirus infection. J. Virol. 70: 7921-7928, 1996. PubMed: <u>8892915</u></p> <p><u>32547</u>: Jang SI, et al . Activator protein 1 activity is involved in the regulation of the cell type-specific expression from the proximal promoter of the human profilaggrin gene. J. Biol. Chem. 271: 24105-24114, 1996. PubMed: <u>8798649</u></p> <p><u>32566</u>: Dittrich E, et al . A di-leucine motif and an upstream serine in the interleukin-6 (IL-6) signal transducer gp130 mediate ligand-induced endocytosis and down-regulation of the IL-6 receptor. J. Biol. Chem. 271: 5487-5494, 1996. PubMed: <u>8621406</u></p> <p><u>32568</u>: Lee JH, et al . The proximal promoter of the human transglutaminase 3 gene. J. Biol. Chem. 271: 4561-4568, 1996. PubMed: <u>8626812</u></p> <p><u>32720</u>: Chen Y, et al . Demonstration of binding of dengue virus envelope protein to target cells. J. Virol. 70: 8765-8772, 1996. PubMed: <u>8971005</u></p> <p><u>32728</u>: Russell DW, Miller AD . Foamy virus vectors. J. Virol. 70: 217-222, 1996. PubMed: <u>8523528</u></p> <p><u>32861</u>: Wright DA, et al . Association of human fas (CD95) with a ubiquitin-conjugating enzyme (UBC-FAP). J. Biol. Chem. 271: 31037-31043, 1996. PubMed: <u>8940097</u></p> <p><u>32893</u>: Zhang J, et al . Dynamin and beta-arrestin reveal distinct mechanisms for G protein-coupled receptor internalization. J. Biol. Chem. 271: 18302-18305, 1996. PubMed: <u>8702465</u></p> <p><u>33011</u>: Ozcelebi F, et al . Phosphorylation of cholecystokinin receptors expressed on chinese hamster ovary cells. J. Biol. Chem. 271: 3750-3755, 1996. PubMed: <u>8631990</u></p> <p><u>33013</u>: Gibson S, et al . Functional LCK is required for optimal CD28-mediated activation of the TEC family tyrosine kinase EMT/ITK. J. Biol. Chem. 271: 7079-7083, 1996. PubMed: <u>8636141</u></p> <p><u>33016</u>: Shaul PW, et al . Acylation targets endothelial nitric-oxide synthase to plasmalemmal caveolae. J. Biol. Chem. 271: 6518-6522, 1996. PubMed: <u>8626455</u></p> <p><u>33032</u>: Ladner RD, et al . Identification of a consensus cyclin-dependent kinase phosphorylation site unique to the nuclear form of human deoxyuridine triphosphate nucleotidohydrolase. J. Biol. Chem. 271: 7752-7757, 1996. PubMed: <u>8631817</u></p> <p><u>33038</u>: Wu X, et al . Demonstration of a physical interaction between microsomal triglyceride transfer protein and apolipoprotein B during the assembly of ApoB-containing lipoproteins. J. Biol. Chem. 271: 10277-10281, 1996. PubMed: <u>8626595</u></p> <p><u>33052</u>: Hawes BE, et al . Phosphatidylinositol 3-kinase is an early intermediate in the G beta gamma-mediated mitogen-activated protein kinase signaling pathway. J. Biol. Chem. 271: 12133-12136, 1996. PubMed: <u>8647803</u></p> <p><u>33137</u>: Arai H, Charo IF . Differential regulation of G-protein-mediated signaling by chemokine receptors. J. Biol. Chem. 271: 21814-21819, 1996. PubMed: <u>8702980</u></p> <p><u>33163</u>: Hsieh CM, et al . APEG-1, a novel gene preferentially expressed in aortic smooth muscle cells, is down-regulated by vascular injury. J. Biol. Chem. 271: 17354-17359, 1996. PubMed: <u>8663449</u></p> <p><u>33175</u>: Holtmann MH, et al . Multiple extracellular loop domains contribute critical determinants for agonist binding and activation of the secretin receptor. J. Biol. Chem. 271: 14944-14949, 1996. PubMed: <u>8663161</u></p>

#### Notices and Disclaimers

ATCC products are intended for laboratory research purposes only. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States and Canada. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your word search in Cell Lines by typing in your search criteria below or returning to the [Cell Lines menu](#). To search another product line, choose one from the dropdown box at the top. For complex searches using boolean operators, the following characters must be used: & (for AND), | (for OR), ^ (for AND NOT). An asterisk (\*) is used as the wildcard. For more information please review the [Search Help](#).

For query options, please read the [search help](#).

#### **ATCC News Archive**

[Home](#) [Ordering Info](#) [Technical Help](#) [About ATCC](#) [Contact Us](#)  
[Privacy Policy](#) [Terms of Use](#) [ATCC MTA](#)

© 2003 American Type Culture Collection (ATCC).  
All rights reserved.